

Exam 09/01/2021 Groups and symmetries

Student

January 2021

1 Theoretical part

1. Given a Lie algebra with generators L^i which have the commutation relations $[L^i, L^j] = i\epsilon^{ijk}L^k$. Is this a real algebra? How can you tell? How can you make this into a real algebra?
2. Describe a root as an element of a dual vector space. Which commutation relation defines these linear maps. These are also solutions of an equation. Which one?
3. Often we work with the components of these roots α^i . In which commutation relations do they appear? How are the components labeled by i defined in the Chevalley basis?
4. The Cartan-Weyl basis defines a real form. How is it called? Is it compact? What is its character?
5. What are the two formal mathematical definitions of a Lie group and what is the relation between them?
6. Suppose a Lie algebra can be split as $\mathfrak{g} = \mathfrak{l} + \mathfrak{h}$. If you impose that \mathfrak{h} is an ideal of \mathfrak{g} what are the commutation relations?
7. Suppose you have a tensor A^{ijk} where $A^{ijk} = -A^{jik}$ and $A^{ijk} + A^{kij} + A^{jki} = 0$. What is the corresponding Young tableau?

2 Practical part

2.1 Question 1: Lorentz algebra

Suppose the Lie algebra $\mathfrak{so}(1,3)$, i.e. Lorentz algebra. It has the generators

$$M_{\mu\nu} = g_{\mu\rho}M^\rho{}_\nu = -M_{\nu\mu} \quad (1)$$

where

$$g_{\mu\nu} = \begin{pmatrix} -1 & 0 \\ 0 & \mathbb{1}_3 \end{pmatrix}; \quad g_{00} = -1; \quad g_{0i} = g_{i0} = 0, \quad g_{ij} = \delta_{ij}. \quad (2)$$

The commutation relations are given by

$$[M_{\mu\nu}, M_{\sigma\rho}] = g_{\nu\sigma}M_{\mu\rho} - g_{\mu\sigma}M_{\nu\rho} - g_{\nu\rho}M_{\mu\sigma} + g_{\mu\rho}M_{\nu\sigma} \quad (3)$$

- (a) There are 6 generators. Can they all 6 be hermitian? Can they all 6 be anti-hermitian?
- (b) Let us define the generator $L^i := \frac{1}{2}\epsilon^{ijk}M_{jk}$. Prove that the commutator is given by $[L^i, L^j] = -\epsilon^{ijk}L^k$.
- (c) Define $K^i := M^i{}_0$. Prove that the following commutators hold

$$[K^i, K^j] = \epsilon^{ijk}L^k; \quad [L^i, K^j] = -\epsilon_{ijk}K^k. \quad (4)$$

- (d) According to the commutation relations, can one realize K^i as anti-hermitian or hermitian generators?

- (e) Calculate the Cartan-Killing metric.
- (f) If we look at the killing form as the trace should K^i be hermitian or anti-hermitian? Explain with the positivity of the trace.
- (g) We have seen that the Lorentz algebra can be realized as a real algebra. Which algebra is this?
- (h) Prove that algebra realized by linear combination of L^i, K^i , i.e. $a_i L^i + b_i K^i$ can also be realized by $(a_i + i b_i) L^i$. Which algebra is this?
- (i) What changes if we set g the euclidean metric, i.e. $g_{00} = 1$?
- (j) Define the algebra $J_{\pm}^i = \frac{1}{2}(L^i \pm K^i)$. What do you find?

2.2 Question 2: Nahm result

Something about bosonic subgroups and fermionic subgroups and supergroups